

国内研究者による文献例

Dynabeads Protein A / Protein G

IP (Immunoprecipitation)

[Nitration and Inactivation of IDO by Peroxynitrite1](#)

Hidetsugu Fujigaki et al

Department of Informative Clinical Medicine, Gifu University Graduate School of Medicine, Gifu City, Japan

[Mass Spectrometry Analysis of the Native Protein Complex Containing Actinin-4 in Prostate Cancer Cells](#)

Tomohiko Hara et al

Chemotherapy Division and Cancer Proteomics Project,
National Cancer Center Research Institute

[Ku70 and Poly\(ADP-Ribose\) Polymerase-1 Competitively Regulate B-Catenin and T-Cell Factor-4–Mediated Gene Transactivation: Possible Linkage of DNA Damage Recognition and Wnt Signaling](#)

Masashi Idogawa et al

Chemotherapy Division and 2ADP-Ribosylation in Oncology Project, National Cancer Center Research Institute, Tokyo, Japan

[Loss of the TOR Kinase Tor2 Mimics Nitrogen Starvation and Activates the Sexual Development Pathway in Fission Yeast](#)

Tomohiko Matsuo et al

Department of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo

[Meichroacidin containing the MORN motif is essential for spermatozoa morphogenesis](#)

Keizo Tokuhira et al

TANAKA Project, Center for Advanced Science and Innovation, Osaka University

[Meiotic association between Spo11 regulated by Rec102, Rec104 and Rec114](#)

Hiroyuki Sasanuma et al

Genetic System Regulation Laboratory, RIKEN Discovery Research Institute

ChIP (Chromatin Immunoprecipitation)

[Escherichia coli Histone-Like Protein H-NS Preferentially Binds to Horizontally Acquired DNA in Association with RNA Polymerase](#)

Taku Oshima et al

Graduate School of Information Science, Nara Institute of Science and Technology

[Reconstruction of the Kinetochores during Meiosis in Fission Yeast](#)

[*Schizosaccharomyces pombe*](#)

Aki Hayashi et al

Kansai Advanced Research Center, National Institute of Information and Communications Technology

[Secondary DNA structure formation for Hoxb9 promoter and identification of its specific binding protein](#)

Takumi Yamagishi et al

Kondo Research Unit, Brain Development Research Group, Brain Science Institute, Institute of Physical and Chemical Research (RIKEN)

[Fission Yeast Taz1 and RPA Are Synergistically Required to Prevent Rapid Telomere Loss](#)

Tatsuya Kibe et al

*Department of Molecular Biotechnology, Graduate School of Advanced Sciences of Matter, Hiroshima University

[Assembly of Regulatory Factors on rRNA and Ribosomal Protein Genes in *Saccharomyces cerevisiae*](#)

Koji Kasahara et al,

Division of Molecular and Cellular Biology, International Graduate School of Arts and Sciences, Yokohama City University

[Interplay between Chromatin and *trans*-Acting Factors on the *IME2* Promoter upon Induction of the Gene at the Onset of Meiosis](#)

Tomomi Inai et al

Department of Molecular Biotechnology, Graduate School of Advanced Sciences of Matter, Hiroshima University,

[Investigation of the Mechanism of Meiotic DNA Cleavage by VMA1-Derived Endonuclease Uncovers a Meiotic Alteration in Chromatin Structure around the Target Site](#)

Tomoyuki Fukuda et al

*Department of Integrated Biosciences, Graduate School of Frontier Sciences,
University of Tokyo*

[Identification of transcriptional regulatory cascades in retinoic acid-induced growth arrest of HepG2 cells](#)

Misato Nakanishi et al

1Laboratory of Genome Exploration Research Group, RIKEN Genomic Sciences Center (GSC),

[Role of Elg1 protein in double strand break repair](#)

Hideaki Ogiwara et al

1Molecular Cell Biology Laboratory, Graduate School of Pharmaceutical Sciences,
Tohoku University