

Dynabeads® Mouse pan B (B220)

Catalog no. 11441D

Store at 2°C to 8°C

Rev. Date: March 2012 (Rev. 008)

Product Contents

Product contents	Volume
Dynabeads® Mouse pan B	5 mL

Product capacity

~2 × 10⁹ cells

Dynabeads® Mouse Pan B contains 4 × 10⁸ beads/mL in phosphate buffered saline (PBS), pH 7.4, containing 0.1% bovine serum albumin (BSA) and 0.02% sodium azide as a preservative.

Caution: Sodium azide may react with lead and copper plumbing to form highly explosive metal azides.

Product Description

This product is intended for positive isolation or depletion of murine B cells directly from spleen, lymph node cell suspensions, or other samples containing B cells. The Dynabeads® are mixed with the cell sample in a tube and will bind to the target cells during a short incubation. The bead-bound cells are separated by a magnet.

Depletion – Discard the bead-bound cells and use the remaining, untouched cells for any application.

Positive isolation – Discard the supernatant and use the bead-bound cells for downstream applications.

Downstream Applications

For rapid and consistent results in protein or gene expression analysis, lyse the B cells while still attached to the beads and directly process for further molecular analysis. For isolation for functional studies, cell activation/expansion, or for flow cytometer analysis, we recommend using Dynabeads® Mouse CD43 to get untouched B cells (bead-free). Dynabeads® are mixed with the cell sample in a tube.

Required Materials

- Magnet (DynaMag™ portfolio). See www.lifetechnologies.com/magnets for recommendations.
- Mixer allowing tilting and rotation of tubes (e.g. HulaMixer® Sample Mixer).
- Isolation Buffer: Ca²⁺ and Mg²⁺ free PBS pH 7.4 with 0.1% (w/v) BSA and 2 mM EDTA.
Note: BSA can be replaced by human serum albumin (HSA) or 2% fetal bovine serum (FBS)/ fetal calf serum (FCS).
- Recommended culture media: RPMI 1640 or DMEM with 10% FCS.

General Guidelines

- Visit www.lifetechnologies.com/samplepreparation for recommended sample preparation procedures.
- Use a mixer that provides tilting and rotation of the tubes to ensure that Dynabeads® do not settle in the tube.
- Keep the temperature at 2°C to 8°C when incubating Dynabeads® and cells, to minimize phagocytic activity and other metabolic processes.
- Follow the recommended volumes and incubation times.
- Avoid air bubbles (foaming) during pipetting.

Protocol

Wash Dynabeads®

See Table 1 for volume recommendations.

1. Resuspend the Dynabeads® in the vial (i.e. vortex for >30 sec, or tilt and rotate for 5 min).
2. Transfer the desired volume of Dynabeads® to a tube.
3. Add the same volume of Isolation Buffer, or at least 1 mL, and resuspend.
4. Place the tube in a magnet for 1 min and discard the supernatant.
5. Remove the tube from the magnet and resuspend the washed Dynabeads® in the same volume of Isolation Buffer as the initial volume transferred of Dynabeads® (step 2).

Prepare Sample

- Prepare a single cell suspension from lymphoid organs (e.g. lymph nodes or spleen) according to “General Guidelines”.
- Resuspend the cells at 1 × 10⁷ cells/mL in Isolation Buffer.

Positively Isolate or Deplete Mouse B Cells

This protocol is based on 1 × 10⁷ cells, but is directly scalable from 1 × 10⁷ to 5 × 10⁸ cells. When working with fewer cells than 1 × 10⁷, use the same volumes as indicated for 1 × 10⁷. When working with higher cell numbers, scale up all volumes accordingly, as shown in Table 1.

1. Transfer 1 mL cells (1 × 10⁷) to a tube and add 25 µL pre-washed and re-suspended Dynabeads®.
2. Incubate for 20 min (positive isolation) or 30 min (depletion) at 2°C to 8°C with gentle tilting and rotation.
3. Place the tube in a magnet for 2 min.
4. For *depletion*; transfer supernatant to a new tube for further use and discard the beads.
or
For *positive isolation*; while the tube is still on the magnet, carefully remove and discard the supernatant.
5. Remove the tube from the magnet and add 1 mL Isolation Buffer, pipet 2–3 times (or vortex 2–3 sec) and place the tube in a magnet for 2 min. While the tube is still in the magnet, carefully remove and discard the supernatant.
6. Repeat step 5 at least once to wash the bead-bound B cells.
This step is critical to obtain a high purity of isolated cells. Resuspend the cell pellet in preferred cell medium.

Keep the cells on 2°C to 8°C until further use in downstream applications.

Table 1: Volumes for isolation/depletion of mouse B cells. This protocol is scalable from 1×10^7 to 5×10^8 cells.

Step	Step description	Volumes per 1×10^7 cells	Volumes per 1×10^8 cells
	Recommended tube size	5 mL	15 mL
	Recommended magnet	DynaMag™-5	DynaMag™-15
1	Cell volume	1 mL	10 mL
1*	Bead volume	25 μ L*	250 μ L*
5-6	For positive isolation only: Wash cells (Isolation Buffer)	$3 \times \sim 1$ mL	$3 \times \sim 10$ mL

* If very high cell-depletion efficiency is required, increase the Dynabeads® volume up to double the recommended amount.

Description of Materials

Dynabeads® Mouse pan B (B220) are uniform, superparamagnetic polystyrene beads (4.5 μ m diameter) coated with a monoclonal antibody specific for the B220 antigen. B220 (also known as CD45R) is expressed on all B cells throughout development, from the early pro-B stage of B cell differentiation.

Related Products

Product	Cat. no.
DynaMag™-5	12303D
DynaMag™-15	12301D
DynaMag™-50	12302D
HulaMixer® Sample Mixer	15920D
Dynabeads® Mouse CD43 (Untouched B Cells)	11422D

REF on labels is the symbol for catalog number.

Limited Use Label License

The purchase of this product conveys to the purchaser the limited, nontransferable right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining additional rights, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008.

Manufactured by Life Technologies AS, Norway. Life Technologies AS complies with the Quality System Standards ISO 9001:2008 and ISO 13485:2003.

Limited Product Warranty

Life Technologies Corporation and/or its affiliate(s) warrant their products as set forth in the Life Technologies' General Terms and Conditions of Sale found on Life Technologies' website at www.lifetechnologies.com/termsandconditions. If you have any questions, please contact Life Technologies at www.lifetechnologies.com/support.

SPEC-05721

©2012 Life Technologies Corporation. All rights reserved. The trademarks mentioned herein are the property of Life Technologies Corporation or their respective owners, except where otherwise stated. LIFE TECHNOLOGIES AND/OR ITS AFFILIATE(S) DISCLAIM ALL WARRANTIES WITH RESPECT TO THIS DOCUMENT, EXPRESSED OR IMPLIED, INCLUDING BUT NOT LIMITED TO THOSE OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. IN NO EVENT SHALL LIFE TECHNOLOGIES AND/OR ITS AFFILIATE(S) BE LIABLE, WHETHER IN CONTRACT, TORT, WARRANTY, OR UNDER ANY STATUTE OR ON ANY OTHER BASIS FOR SPECIAL, INCIDENTAL, INDIRECT, PUNITIVE, MULTIPLE OR CONSEQUENTIAL DAMAGES IN CONNECTION WITH OR ARISING FROM THIS DOCUMENT, INCLUDING BUT NOT LIMITED TO THE USE THEREOF.

For support visit www.lifetechnologies.com/support or email techsupport@lifetech.com

www.lifetechnologies.com

life
technologies™